



























cussed how the hotspot affects the total correlation function. Marking realistic values for plasmonic hotspot sizes and peak amplitudes in our calculations allows the prediction of their benefit for FCS applications. We found that a bow tie antenna can outperform a Mie sphere by two orders of magnitude in terms of increasing the total correlation contrast, due to its field enhancement being  $30\times$  higher than the Mie sphere. Further we showed that for a given hotspot peak intensity, there is an optimal size at which the total correlation contrast becomes maximal. Characteristic parameters of the correlation function like total contrast and diffusion time scale differently with the hotspot width or peak height. This has severe implications: even if the total correlation function seems to be dominated by the hotspot –exhibiting strongly increased contrast or significant reduction of the diffusion time– this does in general not allow to interpret the total correlation function as in a single focus FCS experiment and leads to diffusion coefficients orders of magnitudes different from the real value. We identify hotspot parameter regimes in which an interpretation of the correlation curve as in a single focus experiment is tolerable and yields correct results. Our model is a versatile tool to efficiently predict how non-gaussian intensity distributions alter fluorescence fluctuation measurements. As such it allows to benchmark proposed antenna designs for FCS against the requirements that MDF enhancements of at least around  $10^2$  are obtained and that the antenna sustains such a high MDF over a sufficiently large volume to maximally improve FCS contrast. This is by no means an easy challenge, as large MDF enhancements typically involve narrow gaps.

### **Acknowledgments**

We are grateful to Clara Osorio and Artem Bakulin for carefully reviewing the manuscript. This work is part of the research program of the Foundation for Fundamental Research on Matter (FOM), which is financially supported by the The Netherlands Organization for Scientific Research (NWO). This work is supported by NanoNextNL, a micro- and nanotechnology consortium of the Government of The Netherlands and 130 partners. A.F.K. gratefully acknowledges an NWO-Vidi grant for financial support.